

INDEX TO VOLUME 80(1994)

AUTHORS

- Barnes, J.T. and T.J. Foreman: Recent evidence for the formation of annual growth increments in the otoliths of young Pacific sardines (*Sardinops sagax*), 29-35.
- Barrett, R.H.: see Gogan, 80-83.
- Beamesderfer, R.C.P.: see Rien, 161-170.
- Boroski, B.B. and P. McLaughlin: An efficient technique for capturing swimming deer, 36-42.
- Boydston, L.B.: Analysis of two mark-recapture methods to estimate the fall chinook salmon (*Oncorhynchus tshawytscha*) spawning run in Bogus Creek, California, 1-13.
- Cech, J.J. Jr.: see Setka, 89-96.
- Cordone, A.: In memoriam: Leo Shapovalov, 181.
- Davis, L.H.: Review of: Wild pigs of the United States: their history, morphology, and current status, 129-130.
- Fleskes, J.P., J.M. Hicks, D.S. Gilmer, and D. Yparraguirre: Changing patterns of goose harvest on California public hunting areas, 133-149.
- Foreman, T.J.: see Barnes, 29-35.
- Foster, C.A.: see Rien, 161-170.
- Garrison, B.A.: Review of: California forests and woodlands: a natural history, 128-129.
- Garrison, B.A.: Determining the biological significance of changes in predicted habitat values from the California Wildlife Habitat Relationships System, 150-160.
- Gilmer, D.S.: see Fleskes, 133-149.
- Gobalet, K.W.: A prehistoric sturgeon fishery in San Pablo, Contra Costa County, California: an addendum, 125-127.
- Gogan, P.J. and R.H. Barrett, Roosevelt elk dietary quality in northern coastal California, 80-83.
- Hicks, J.M.: see Fleskes, 133-149.
- Hothem, R.L. and D. Welsh: Duck and shorebird reproduction in the grasslands of central California, 68-79.
- Kalvass, P.E.: The effect of different harvest methods on sea palm (*Postelsia palmaeformis*) sporophyll growth, 57-67.
- Law, P.M.W.: Simulation study of salmon carcass survey by capture-recapture methods, 14-28.
- Leos, R.R.: A portable field sampling table for dockside sampling of fish, 84-87.
- McLaughlin, P.: see Boroski, 36-42.
- Rien, T.A., R.C.P. Beamesderfer, and C.A. Foster: Retention, recognition, and effects on survival of several tags and marks for white sturgeon, 161-170.
- Setka, J.D. and J.J. Cech, Jr.: Blood and muscle characteristics of leopard shark (*Triakis semifasciata*) and brown smoothhound (*Mustelus henlei*), 89-96.
- Wendell, F.: Relationship between sea otter range extension and red abalone abundance and size distribution in central California, 45-56.
- Welsh, D.: see Hothem, 68-79.
- Welsh, H.H. Jr.: Bioregions: an ecological and evolutionary perspective and a proposal for California, 97-124.
- Whitmore, D. Waterfowl killed by lightning, 43-44.
- Wooster, T. An observation of high fecundity of spotted owls in California, 171.
- Yparraguirre, D.: see Fleskes, 133-149.

INDEX TO VOLUME 80(1994)

AUTHORS

- Barnes, J.T. and T.J. Foreman: Recent evidence for the formation of annual growth increments in the otoliths of young Pacific sardines (*Sardinops sagax*), 29-35.
- Barrett, R.H.: see Gogan, 80-83.
- Beamesderfer, R.C.P.: see Rien, 161-170.
- Boroski, B.B. and P. McLaughlin: An efficient technique for capturing swimming deer, 36-42.
- Boydston, L.B.: Analysis of two mark-recapture methods to estimate the fall chinook salmon (*Oncorhynchus tshawytscha*) spawning run in Bogus Creek, California, 1-13.
- Cech, J.J. Jr.: see Setka, 89-96.
- Cordone, A.: In memoriam: Leo Shapovalov, 181.
- Davis, L.H.: Review of: Wild pigs of the United States: their history, morphology, and current status, 129-130.
- Fleskes, J.P., J.M. Hicks, D.S. Gilmer, and D. Yparraguirre: Changing patterns of goose harvest on California public hunting areas, 133-149.
- Foreman, T.J.: see Barnes, 29-35.
- Foster, C.A.: see Rien, 161-170.
- Garrison, B.A.: Review of: California forests and woodlands: a natural history, 128-129.
- Garrison, B.A.: Determining the biological significance of changes in predicted habitat values from the California Wildlife Habitat Relationships System, 150-160.
- Gilmer, D.S.: see Fleskes, 133-149.
- Gobalet, K.W.: A prehistoric sturgeon fishery in San Pablo, Contra Costa County, California: an addendum, 125-127.
- Gogan, P.J. and R.H. Barrett, Roosevelt elk dietary quality in northern coastal California, 80-83.
- Hicks, J.M.: see Fleskes, 133-149.
- Hothem, R.L. and D. Welsh: Duck and shorebird reproduction in the grasslands of central California, 68-79.
- Kalvass, P.E.: The effect of different harvest methods on sea palm (*Postelsia palmaeformis*) sporophyll growth, 57-67.
- Law, P.M.W.: Simulation study of salmon carcass survey by capture-recapture methods, 14-28.
- Leos, R.R.: A portable field sampling table for dockside sampling of fish, 84-87.
- McLaughlin, P.: see Boroski, 36-42.
- Rien, T.A., R.C.P. Beamesderfer, and C.A. Foster: Retention, recognition, and effects on survival of several tags and marks for white sturgeon, 161-170.
- Setka, J.D. and J.J. Cech, Jr.: Blood and muscle characteristics of leopard shark (*Triakis semifasciata*) and brown smoothhound (*Mustelus henlei*), 89-96.
- Wendell, F.: Relationship between sea otter range extension and red abalone abundance and size distribution in central California, 45-56.
- Welsh, D.: see Hothem, 68-79.
- Welsh, H.H. Jr.: Bioregions: an ecological and evolutionary perspective and a proposal for California, 97-124.
- Whitmore, D. Waterfowl killed by lightning, 43-44.
- Wooster, T. An observation of high fecundity of spotted owls in California, 171.
- Yparraguirre, D.: see Fleskes, 133-149.

SUBJECT

- Abalone, Red: Abundance in relation to sea otter, 45-56.
Bioregions: Proposal for California, 97-124.
Deer, Black-tailed: Capturing swimming deer, 36-42.
Ducks: Reproduction in central California grasslands, 68-79.
Elk, Roosevelt: Dietary quality, 80-83.
Geese: Patterns of goose harvest, 133-149.
Habitat, Wildlife: Significance of change in habitat value, 150-160.
Lightning: Waterfowl killed by lightning, 43-44.
Mark-recapture: Analysis of chinook salmon populations, 1-13.
Mark-recapture: Simulation of salmon carcass survey, 14-28.
Palm, Sea: Effect of harvest methods, 57-67.
Otter, Sea: Range expansion, 45-56.
Owl, Spotted: Observation of high fecundity, 171.
Salmon, Chinook: Mark-recapture methods, 1-13, 14-28.
Sampling, Field: Portable table, 84-87.
Sardine, Pacific: Annual growth, 29-35.
Shark, Leopard: Blood and muscle characteristics, 89-96.
Shorebirds: Reproduction in central California grasslands, 68-79.
Smoothhound, Brown: Blood and muscle characteristics, 89-96.
Sturgeon: Prehistoric fishery in San Pablo, 125-127.
Sturgeon, White: Tag and marking retention and recongition, 161-170.
Waterfowl: Lightning kill, 43-44.

SPECIES

- Acipenser medirostris*, 125-127.
Acipenser trasmontanus, 125-127, 161-170.
Anas acuta, 43-44, 68-79.
Anas cyanoptera, 68-79.
Anas platyrhynchos, 68-79.
Anas strepera, 68-79.
Anser albifrons frontalis, 133-149.
Anser caeruleus caeruleus, 43-44.
Anser rossii, 43-44.
Branta canadensis minima, 133-149.
Cervus elaphus roosevelti, 80-83.
Charadrius vociferus, 68-79.
Chen caeruleus caeruleus, 133-149.
Chen rossi, 133-149.
Enhydra lutra, 45-56.
Haliotis rufescens, 45-56.
Himantopus mexicanus, 68-79.
Mustelus henlei, 89-96.
Odocoileus hemionus columbianus, 36-42.
Oncorhynchus sp., 14-28, 125-127.
Oncorhynchus tshawytscha, 1-13.

- Postelsia palmaeformis*, 57-67.
Recurvirostra americana, 68-79.
Sardinops sagax, 29-35.
Strix occidentalis caurina, 171
Triakis semifasciata, 89-96.

